

0.05% Trypsin-EDTA in PBS (500mL; TBS8085)

DESCRIPTION

Trypsin-EDTA solutions are commonly used to detach adherent cells from culture surfaces. Various concentrations of trypsin-EDTA are available, and their use depends primarily on the cell type and the age of the culture to dislodge cells from the substrate. The general recommended working concentration is 0.05% Trypsin for cell detachment.

0.05% Trypsin-EDTA is a ready-to-use solution (1x) without additional dilution.

COMPONENTS:

Trypsin: 0.05%
EDTA: 1mM
Phenol Red: 10mg/L
1x PBS
pH: 7.2-80

PACK SIZE

500 mL /bottle, Sterile.
Store at -20°C.

PREPERATION FOR CELL CULTURE

This product is recommended to be directly used for cell detachment without additional dilution.

RELATED PRODUCTS

MSC Medium (TBS8021)
Ham's F-12 Medium (TBS8032)
DMEM Medium (TBS6061)
Chondrogenic Differentiation Medium (TBS8062)
RPMI16040 Medium (TBS8063)
Hybridoma Growth Medium (TBS8074)
DMEM-F12 Mixture (TBS8083)
ESC/iPSC-qualified FBS (TBS8002)
Adipocyte Differentiation Cocktail (TBS8017)
0.1% Gelatin Solution (TBS8004)
2x HBS, pH7.05 (TBS5076)
Cell Culture Grad Water (TBS5050)
LB Medium (TBS8056)
HAT Supplement (TBS8075)
HT Supplement (TBS8073)
Hybridoma Growth Medium (TBS8074)
Pen-strep-fug solution (TBS8037)
Pen-strep solution (TBS8029)

Research use only.