

Glucose-6-phosphate dehydrogenase, Enzyme Activity

Catalog	Unit
TBP0017-1KU	1000 U
TBP0017-5KU	5000 U

Preparation and Specification

Appearance: White amorphous powder, lyophilized

Activity: Gradelll 200U/mg-solid or more

Contaminants: Creatine phosphokinase $\leq 1 \times 10^{-3}\%$

Phosphoglucomutase $\leq 1 \times 10^{-3}\%$

6-Phosphogluconate dehydrogenase $\leq 5 \times 10^{-3}\%$

Phosphoglucose isomerase $\leq 1 \times 10^{-2}\%$

Glutathione reductase $\leq 1 \times 10^{-3}\%$


Hexokinase $\leq 1 \times 10^{-2}\%$

Myokinase $\leq 1 \times 10^{-2}\%$

NADH oxidase $\leq 1 \times 10^{-2}\%$

NADPH oxidase $\leq 1 \times 10^{-2}\%$

D-Glucose-6-phosphate:NADP⁺ 1-oxidoreductase (EC 1.1.1.49)



Properties

Stability: Stable at -20°C for at least One year

Molecular weight: approx. 140,000 (by gel filtration)

Michaelis constants: NAD⁺ linked 2.4×10^{-4} M (NAD⁺), 4.7×10^{-4} M (G-6-P)

NADP⁺ linked 7.4×10^{-6} M (NADP⁺), 3.2×10^{-4} M (G-6-P)

Inhibitors: Metal ions, iodoacetamimide, SDS etc.

Optimum pH: 7.8

Optimum temperature: 50°C-55°C

pH Stability: pH 5.0-11.0 (25°C, 22hr)

Thermal stability: below 50°C (pH 7.8, 30min)

Applications

The enzyme is useful for enzymatic determination of NAD⁺(NADP⁺) and G-6-P, and activities of phosphoglucose isomerase, phosphoglucomutase and hexokinase. The enzyme is also used for enzymatic determination of glucose and creatine phosphokinase activity when coupled with hexokinase (HXK-311).

For research use only