

## Phosphoenolpyruvate carboxylase, Enzyme Activity

Catalog	Unit
TBP0039-1KU	1000 U
TBP0039-5KU	5000 U

## **Preparation and Specification**

Appearance: White amorphous powder, lyophilized

Activity: GradeIII 5.0U/mg-solid or more

Contaminants: Lactate dehydrogenase  $\leq 1.0 \times 10^{-3}\%$ 

Pyruvate kinase ≤0.5%

Stabilizers: BSA, sugar alcohols

Orthophosphate: oxaloacetate carboxy-lyase (Phosphorylating) (EC 4.1.1.31)

Phosphoenolpyruvate+CO2+H2O-Oxaloacetate+Pi Mg<sup>++</sup>

## **Properties**

Stability: Stable at -20°C for at least One year

Molecular weight: approx. 390,000 (by gel filtration)

Isoelectric point: 6.0±0.1

Structure: 4 Subunits (M.W.100,000) per enzyme molecule

Michaelis constant: 1.9×10<sup>-4</sup>M (Phosphoenolpyruvate)

Optimum pH: 7.5-8.0

Optimum temperature: 60°C

pH Stability: pH 5.0-8.0 (25°C, 24hr)

Thermal stability: below 40°C (pH 7.0, 15min)

## **Applications**

This enzyme is useful for enzymatic determination of carbon dioxide when coupled with malate dehydrogenase (MAD-201, MAD-211) in clinical analysis.

For research use only