

Catalog	Kit Size
TBS32121-48	48 assays
TBS32121-96	96 assays

INTRODUCTION

Acetylation of histone H3 at lysine 9 (H3K9) is a post-translational modification where an acetyl group is added to the lysine residue at position 9 on the histone H3 protein. This modification is mediated by histone acetyltransferases (HATs), which add acetyl groups, and histone deacetylases (HDACs), which remove them. H3K9 acetylation is associated with gene activation. It is often found at active promoters and enhancers, facilitating a more open chromatin structure that allows transcription factors to access DNA. Altered acetylation levels at H3K9 have been linked to various cancers, affecting tumor progression and metastasis.

The Global Acetyl Histone H3K9 Quantification Kit is designed to measure global histone H3K9 acetylation in various mammalian cells, including human and mouse cells, as well as fresh and frozen tissues, cultured adherent cells, and suspension cells. This assay is simple, user-friendly, accurate, and fast. The measurement can be completed in 2.5 hours, with a detection range of 20 ng to 5 µg per well.

Alternative Name: H3K9ac; Acetyl Histone H3K9; Acetylation of histone H3 at lysine 9

PRINCIPLE OF THE ASSAY

In this assay, an acetylated histone H3 at lysine 9 binds to the capture antibody against a specific acetylated histone H3K9 which is immobilized within the strip wells to make an antibody-antigen complex. Following this step, the bound acetyl histone H3K9 is detected using a secondary antibody conjugated to a detectable label. A subsequent reaction with a color development reagent produces a measurable signal, where the intensity of absorbance is directly proportional to the relative amount of acetylated histone H3K9 present in the sample. The absolute amount of acetyl histone H3K9 is quantified from a control standard with known concentrations, ensuring precise measurement.

KIT CONTENT AND STORAGE CONDITIONS

PART	48 ASSAYS	96 ASSAYS	STORAGE OF OPENED/ RECONSTITUTED
Detection Antibody	5 µL of histone H3K9 antibody(1000x).	10 µL of histone H3K9 antibody(1000x).	-20°C
Histone H3K9 Standard	10 µL of histone H3K9 control.	20 µL of histone H3K9 control.	-20°C
Stop Solution	3 mL of 2 N sulfuric acid.	6 mL of 2 N sulfuric acid.	Room temperature away from light
Wash Buffer	10 mL of concentrated solution (10x).	20 mL of concentrated solution (10x).	May be stored for up to 6 months at 2-8 °C.
Antibody Buffer	6 ml	12 ml	
Color Developer	5 ml	10 ml	
8-Well Sample Strips (with Frame) Precoated with H3K9 antibody	6	12	

Store the unopened kit at 2-8 °C. Do not use past kit expiration date.

PRECAUTIONS

Wear protective gloves, clothing, eyes, and face protection. Wash hands thoroughly after handling.


PROTOCOL

- Determine the number of strip wells required. Leave these strips in the plate frame (remaining unused strips can be placed back in the bag. Seal the bag tightly and store at 4°C).
- Dilute 10x **Wash Buffer** with distilled water at a 1:9 ratio (ex: 1 ml of 10x **Wash Buffer** + 9 ml of water) to 1x Wash Buffer.
- Add 50 µl of **Sample** into sample well (sample protein amount: 50-200 ng of the histone extract).
- Standard Preparation: Label test tubes as #1 through #8. Pipet 294 µL of 1x Antibody Buffer into tube #1, and 150 µL into tubes #2 to #7 **as diagram below (Fig.2)**.

Add 6 μL of the Standard Control (100 ng/μL) to tube #1, and mix. Then, make 2x serial dilutions of the standard using the tube #1 standard solution from tube #2 through #7 with sequential transfer of 150 μL to the next concentration. Mix each tube thoroughly before the next transfer.

Add 50 μL of each standard concentration into the indicated standard well. The standard amount of 50 μL in tubes 1 through 7 will be 100, 50, 25, 12.5, 6.25, 3.13, and 1.56 ng/well. Tube# 8 is Standard 0.

Fig.2 Diagram for Acetyl Histone H3K9 standard preparation



- Aspirate and wash the wells with 150 μl of 1x **Wash Buffer** three times.
- Dilute **Detection Antibody** (at a 1:1000 ratio) to working detection with **Antibody Buffer**. Add 50 μl of diluted **Detection Antibody** to each well and incubate at room temperature for 60 minutes on an orbital shaker (100 rpm).
- Aspirate and wash the wells with 150 μl of 1x **Wash Buffer** for 3x times.
- Add 100 μl of **Color Developer** into the wells and incubate at room temperature for 2-10 minutes away from light. Monitor the color development in the sample and standard wells (blue).
- Add 50 μl of **Stop Solution** to each well to stop enzyme reaction when the color in the standard wells containing the higher concentrations of standard control turns medium blue. The color should change to yellow and absorbance can be read on a microplate reader at 450 nm within 2-15 minutes.
- Calculate % histone H3K9 acetylation:

$$Acetylation \% = \frac{OD (treated (tested) sample - blank)}{OD (untreated (control) sample - blank)} \times 100\%$$

For the amount quantification, plot OD versus amount of Standard Control and determine the slope as delta OD/ng.

Calculate the amount of acetyl H3K9 using the following formula:

$$Amount (ng/mg protein) = \frac{OD (sample - blank)}{Protein (\mu g) * slope} \times 1000$$

* Histone extract amount added into the sample well at step 3.

HISTONE EXTRACTION PROTOCOL

1. For tissues (treated and untreated), weigh the sample and cut the sample into small pieces (1-2 mm³) with a scalpel or scissors. Transfer tissue pieces to a Dounce homogenizer. Add TEB buffer or PBS containing 0.5% Triton X 100 or 2 mM PMSF and 0.02% NaN₃ at 200 mg/ml, and disaggregate tissue pieces by 50-60 strokes. Transfer homogenized mixture to a 15 ml conical tube and centrifuge at 3000 rpm for 5 minutes at 4°C. If total mixture volume is less than 2 ml, transfer mixture to a 2 ml vial and centrifuge at 10,000 rpm for 1 minute at 4°C. Remove supernatant.

For cells (treated and untreated), harvest cells and pellet the cells by centrifugation at 1000 rpm for 5 minutes at 4°C. Resuspend cells in TEB buffer at 10⁷ cells/ml and lyse cells on ice for 10 minutes with gentle stirring. Centrifuge at 3000 rpm for 5 minutes at 4°C. If total volume is less than 2 ml, transfer cell lysates to a 2 ml vial and centrifuge at 10,000 rpm for 1 minute at 4°C. Remove supernatant.

2. Resuspend cell/tissue pellet in 3 volumes (approx. 200 µl/10⁷ cells or 200 mg of tissue) of extraction buffer (0.5N HCl + 10% glycerol) and incubate on ice for 30 minutes.
3. Centrifuge at 12,000 rpm for 5 minutes at 4°C and remove the supernatant fraction to a new vial.
4. Add 8 volumes (approx. 0.6 ml/10⁷ cells or 200 mg of tissue) of acetone and leave at -20°C overnight.
5. Centrifuge at 12,000 rpm for 5 minutes and air-dry the pellet. Dissolve the pellet in distilled water (30-50 µl/10⁷ cells or 200 mg of tissue).
6. Quantify the protein concentration. Aliquot the extract and store the extract at -20°C or -80°C.

RELATIVE PRODUCTS

Total Histone H3 Acetylation Detection (Colorimetric) (TBS32120)

Total Histone H3 Acetylation Detection (Fluorometric) (TBS32122)

Global Histone H3K4 Methylation Assay Kit (TBS32123)

Global Tri-Methyl Histone H3K4 Quantification Kit (Colorimetric) (TBS32124)

Histone H3 Modification Multiplex Assay Kit (Colorimetric) (TBS32125)

For research use only.